



Radiation Dose in X-Ray and CT Exams

What are x-rays and what do they do?

X-rays are a form of energy, similar to light and radio waves. X-rays are also called radiation. Unlike light waves, x-rays have enough energy to pass through your body. As the radiation moves through your body, it passes through bones, tissues and organs differently, which allows a radiologist to create images of them. The radiologist is a specially trained physician who can examine these images on a monitor. The monitor is like a computer display. It allows the radiologist to see very fine detail of the structures in your body.

X-ray examinations provide valuable information about your health and help your doctor make an accurate diagnosis. X-rays are sometimes used to help place tubes or other devices in the body or to treat disease.



See "Safety in X-ray, Interventional Radiology and Nuclear Medicine Procedures" for more information.

Measuring radiation dosage

When radiation passes through the body, some of it gets absorbed. The x-rays that are not absorbed are used to create the image. The amount that is absorbed contributes to the patient's radiation dose. The radiation that passes through the body does not. The scientific unit of measurement for whole body radiation dose, called "effective dose," is the millisievert (mSv). Other radiation dose measurement units include rad, rem, roentgen, sievert, and gray.

Doctors use "effective dose" when they talk about the risk of radiation to the entire body. Risk refers to

possible side effects, such as the chance of developing a cancer later in life. Effective dose takes into account how sensitive different tissues are to radiation. If you have an x-ray exam that includes tissues or organs that are more sensitive to radiation, your effective dose will be higher. Effective dose allows your doctor to evaluate your risk and compare it to common, everyday sources of exposure, such as natural background radiation.

Naturally-occurring "background" radiation


We are exposed to natural sources of radiation all the time. According to recent estimates, the average person in the U.S. receives an effective dose of about 3 mSv per year from natural radiation, which includes cosmic radiation from outer space. These natural "background doses" vary according to where you live.

People living at high altitudes such as Colorado or New Mexico receive about 1.5 mSv more per year than those living near sea level. A coast-to-coast round trip airline flight is about 0.03 mSv due to exposure to cosmic rays. The largest source of background radiation comes from radon gas in our homes (about 2 mSv per year). Like other sources of background radiation, the amount of radon exposure varies widely depending on where you live.

To put it simply, the amount of radiation from one adult chest x-ray (0.1 mSv) is about the same as 10 days of natural background radiation that we are all exposed to as part of our daily living.

Effective radiation dose in adults

Here are some approximate comparisons of background radiation and effective radiation dose in adults for several radiology procedures described on this website.

	Procedure	Approximate effective radiation dose	Comparable to natural background radiation for:
ABDOMINAL REGION 	Computed Tomography (CT)–Abdomen and Pelvis	10 mSv	3 years
	Computed Tomography (CT)–Abdomen and Pelvis, repeated with and without contrast material	20 mSv	7 years
	Computed Tomography (CT)–Colonography	6 mSv	2 years
	Intravenous Pyelogram (IVP)	3 mSv	1 year
	Barium Enema (Lower GI X-ray)	8 mSv	3 years
	Upper GI Study with Barium	6 mSv	2 years

BONE**Procedure****Approximate effective radiation dose****Comparable to natural background radiation for:**

Spine X-ray

1.5 mSv

6 months

Extremity (hand, foot, etc.) X-ray

0.001 mSv

3 hours

CENTRAL NERVOUS SYSTEM**Procedure****Approximate effective radiation dose****Comparable to natural background radiation for:**

Computed Tomography (CT)–Head

2 mSv

8 months

Computed Tomography (CT)–Head, repeated with and without contrast material

4 mSv

16 months

Computed Tomography (CT)–Spine

6 mSv

2 years

CHEST**Procedure****Approximate effective radiation dose****Comparable to natural background radiation for:**

Computed Tomography (CT)–Chest

7 mSv

2 years

Computed Tomography (CT)–Lung Cancer Screening

1.5 mSv

6 months

Chest X-ray

0.1 mSv

10 days

DENTAL**Procedure****Approximate effective radiation dose****Comparable to natural background radiation for:**

Dental X-ray

0.005 mSv

1 day

HEART**Procedure****Approximate effective radiation dose****Comparable to natural background radiation for:**

Coronary Computed Tomography Angiography (CTA)

12 mSv

4 years

Cardiac CT for Calcium Scoring

3 mSv

1 year

MEN'S IMAGING



Procedure

Approximate
effective
radiation dose


Comparable to
natural
background
radiation for:

Bone Densitometry (DEXA)

0.001 mSv

3 hours

NUCLEAR MEDICINE



Procedure

Approximate
effective
radiation dose

Comparable to
natural
background
radiation for:

Positron Emission
Tomography–Computed Tomography
(PET/CT)

25 mSv

8 years

WOMEN'S IMAGING



Procedure

Approximate
effective
radiation dose

Comparable to
natural
background
radiation for:

Bone Densitometry (DEXA)

0.001 mSv

3 hours

Mammography

0.4 mSv

7 weeks

Note for pediatric patients: Pediatric patients vary in size. Doses given to pediatric patients will vary significantly from those given to adults. For more information on radiation safety in pediatric imaging, visit <http://www.imagegently.org/Roles-What-can-I-do/Parent>.

* The effective doses are typical values for an average-sized adult. The actual dose can vary substantially, depending on a person's size as well as on differences in imaging practices.

Please note that this chart attempts to simplify a very complex topic. If you have questions about radiation risk, talk to your medical physicist and/or radiologist and ask about the benefits and risks of radiologic care.

The International Commission on Radiological Protection (ICRP) Report 103 states: "The use of effective dose for assessing the exposure of patients has severe limitations that must be considered when quantifying medical exposure," and "The assessment and interpretation of effective dose from medical exposure of patients is very problematic when organs and tissues receive only partial exposure or a very heterogeneous exposure which is the case especially with x-ray diagnostics." In other words, effective dose is not always the same for everyone. It can vary based on a person's height and weight, how the procedure is performed and the area of the body being exposed to radiation.

Benefit versus risk

The risk associated with medical imaging procedures refers to possible long-term or short-term side effects. Most imaging procedures have a relatively low risk. Hospitals and imaging centers apply the principles of ALARA (As Low As Reasonably Achievable). This means they make every effort to decrease radiation risk. It is important to remember that a person is at risk if the doctor cannot accurately diagnose an illness or injury. Therefore, it could be said that the benefit from medical imaging, which is an accurate diagnosis, is greater than the small risk that comes with using it. Talk to your doctor or radiologist about any concerns you may have about the risks of a procedure.

For more discussions about benefit versus risk, see the Benefits and Risks section

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2019 Radiological Society of North America, Inc.